

Live Well to
Learn Well

Web:
health.cornell.edu

Phone (24/7):
607-255-5155

Fax:
607-255-0269

Appointments:
Monday–Saturday

*Check web for hours,
services, providers,
and appointment
information*

110 Ho Plaza,
Ithaca, NY
14853-3101

A concussion is an injury to the brain usually caused by a blow to the head. In most cases, it does not involve a loss of consciousness. When you've had a head injury, it's important to know:

- the symptoms of a concussion
- how it should be managed
- its lasting effects

Monitor your symptoms

Because the brain is very complex, every brain injury is different. Some symptoms may appear right away; others may not show up for days after the concussion.

Seek medical care if you experience any of the following symptoms after suffering from a head injury:

- headaches that won't go away
- having trouble (more than usual):
 - remembering things
 - paying attention or concentrating
 - organizing daily tasks
 - making decisions or solving problems
- slowness in thinking, acting, speaking, or reading
- getting lost or easily confused
- neck pain
- feeling tired all the time, lack of energy
- change in sleeping pattern
 - sleeping for much longer periods of time than before
 - trouble sleeping or insomnia
- loss of balance, feeling dizzy, light-headed
- increased sensitivity to light, sound, distractions
- blurred vision, eyes that tire easily
- ringing in the ears
- mood changes:
 - feeling sad, anxious, listless
 - becoming easily irritated or angry for little or no reason
 - lack of motivation

Sometimes the injury makes it hard for people to recognize or acknowledge they are having problems. It may be helpful to enlist the support of others for help with recognizing symptoms. Most individuals can be safely observed at home by friends, roommates, or family members. In fact, friends and family members may be better at noticing changes in behavior than a medical provider (who may not know the patient as well).

Symptoms of a concussion are usually temporary, but may last for days or weeks. Watch for new symptoms. If you feel something is "not quite right," or if you're "feeling foggy," talk with your health care provider.



Rest & careful observation are needed after concussion.

RECOGNIZE DANGER SIGNS

Visit your health care provider or emergency department immediately if —after a blow to the head—you have any of these signs:

- headaches that get worse
- weak or numb arms or legs
- unsteadiness
- repeated vomiting

The people checking on you should take you to the emergency department right away if you:


- are very drowsy or cannot be awakened
- cannot recognize people or places
- have one pupil that is larger than the other
- have convulsions or seizures
- have slurred speech
- behave unusually, seem confused or are very irritable

Facilitate full recovery

While signs of concussion can last for weeks, most people recover without any long-term problems. The two most important things you can do to facilitate a full recovery are rest and careful observation. Resting your mind and body can be hard for busy, highly motivated students; however, taking shortcuts may compromise your ability to return to work and play at pre-concussion levels.

Take care of your body

You should not return to strenuous activity (including sports) while you still have signs of a concussion, like headache or dizziness. Other suggestions include:



- Get plenty of sleep.
- Ask your health care provider when you are ready to drive a car, ride a bike, etc.
- Don't drink alcoholic beverages until your health care provider says you are well enough to do so. Alcohol and certain other drugs may slow your recovery and put you at risk of further injury.
- Take only drugs approved by your health care provider. Avoid any medications that cause drowsiness or changes in level of consciousness: pain medications, sleeping pills, muscle relaxants, tranquilizers, or recreational drugs.

Take care of your brain

Reducing cognitive strain during the healing process is critical to allowing the brain to heal. You'll need to be patient because healing takes time. Prematurely engaging in activities involving mental exertion or multitasking (e.g., reading, taking notes, problem-solving, test-taking, computer or smartphone use) may exacerbate or prolong symptoms and decrease your ability to perform at your pre-concussion level.

- Return to daily activities, such as work or school, gradually and at your own pace.
- Complete the Student Disability Services self-disclosure form.
- Reach out to professors and meet regularly with your academic advising office to discuss your symptoms and develop a plan for return to study.
- If you need additional assistance, speak with your primary care provider.
- Write things down if you are having trouble remembering.
- If you're easily distracted, try to do just one thing at a time (e.g., don't watch TV while preparing dinner).
- Consult your family members or close friends if making important decisions.

Don't risk 2nd brain injury

During the healing process, you should be extremely careful to avoid activities that could lead to another blow or a jolt to your head (including contact or recreational sports). Suffering a second brain injury can cause a much longer recovery.

Recovery time

Most people recover from a first concussion in 7-10 days. How quickly

they improve depends on a variety of factors, including:

- the amount and severity of symptoms
- what part of the brain was injured
- their age
- how healthy they were before the concussion

If you already had a medical problem at the time of your concussion, it may take longer for you to recover from your brain injury. Anxiety and depression may also make it harder to adjust to the symptoms of brain injury.

Be sure to schedule follow-up appointments with your health care provider as recommended. If symptoms increase or do not improve within 24 hours of consulting with a health care provider, call Cornell Health (607-255-5155) for follow-up care right away. If symptoms persist beyond two weeks, contact Cornell Health to discuss a referral to Student Disability Services for appropriate academic accommodations.

For more information

- [cdc.gov/Concussion](https://www.cdc.gov/concussion/)
- [imPACTtest.com](https://www.imPACTtest.com)
- [ncaa.org](https://www.ncaa.org) [search "concussion"]

FOR INTERCOLLEGIATE ATHLETES

Return to play

Report all head injuries or symptoms of concussion to your team athletic trainer immediately. Never return to your sport or a heavy activity level after a major concussion until you've gone through the following stages of recovery and been cleared by your team physician:

Stages of recovery

- Physical & mental rest for 24-48 hours
- Light aerobic exercise (e.g., walking, stationary cycling), no resistance training
- Sport-specific exercise (e.g., skating or running drills), no head impact activities
- Non-contact training drills (start light, progressive resistance training)
- Full contact practice after clearance by team physician & athletic trainer
- Return to competition

At least 24 hours should be devoted to each stage of recovery. If symptoms worsen, stop and resume activities the following day. Resistance training should be added only in the later stages.

ImPACT & SCAT 5 testing

All Cornell varsity athletes will have baseline neuropsychological (imPACT) testing and baseline SCAT 5 test before starting their sport. Post-injury testing will be used as a tool to aid in returning the athlete to their sport. The SCAT 5 test includes balance testing to assess the vestibular system* which can be negatively affected by concussion.

** A system in the body that controls sense of movement and balance.*



Return to Learn

- Complete the Student Disability Services self-disclosure form.
- If you need additional assistance, reach out to your athletic trainer or team physician.